

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Agriculture and Natural Resources

1-16-2016

SCN Resistance Continues to Pay Twice in 2015

Gregory L. Tylka

Iowa State University, gltylka@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Plant Pathology Commons](#)

Recommended Citation

Tylka, Gregory L., "SCN Resistance Continues to Pay Twice in 2015" (2016). *Integrated Crop Management News*. 2325.
<http://lib.dr.iastate.edu/cropnews/2325>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

SCN Resistance Continues to Pay Twice in 2015

Abstract

Resistant soybean varieties are a vital tool for managing the soybean cyst nematode (SCN). These soybean varieties must contain several genes to be fully resistant to the nematode. Not all SCN-resistant varieties contain a full set of resistance genes. Consequently, SCN-resistant soybean varieties can vary greatly in their agronomic performance and the SCN control that they provide.

Keywords

Plant Pathology

Disciplines

Agricultural Science | Agriculture | Plant Pathology

IOWA STATE UNIVERSITY

Extension and Outreach

Integrated Crop Management

[Home](#)

Mailing Lists

Subscribe to ICM News updates and receive email alerts when new information is posted.

Your Email address *

SCN Resistance Continues to Pay Twice in 2015

ICM News

January 16, 2016

Resistant soybean varieties are a vital tool for managing the soybean cyst nematode (SCN). These soybean varieties must contain several genes to be fully resistant to the nematode. Not all SCN-resistant varieties contain a full set of resistance genes. Consequently, SCN-resistant soybean varieties can vary greatly in their agronomic performance and the SCN control that they provide.

ISU SCN-resistant Soybean Variety Trial Program

ISU evaluates the performance of more than 150 SCN-resistant soybean varieties each year. The research is supported by soybean checkoff funding through a grant from the Iowa Soybean Association. Experiments are conducted in nine SCN-infested fields in northern, central, and southern Iowa (figure 1). Soybean yield and beginning- and end-of-season SCN soil egg population densities (numbers) are measured from each of the thousands of individual 4-row-wide plots (figure 2). Several susceptible (non resistant)

soybean varieties are included in each experiment for comparison. Most of the SCN-resistant soybean varieties in the experiments are glyphosate resistant, but some conventional varieties with no herbicide resistance also are evaluated in the experiments.




Figure 1. Locations of the nine SCN-resistant Soybean Variety Trial Program field experiments in 2015.




Figure 2. Aerial view of SCN-resistant Soybean Variety Trial Program field experiment and other research plots in east central Iowa, near Urbana, in 2015.

Processing of the SCN soil samples collected at harvest was completed in December, and the final report of the experimental results has been compiled. A printed copy of the report was distributed as a special insert in the January 16, 2016, issue of Iowa Farmer Today. The report also is available electronically at www.isuscntrials.info and for free download at the Iowa State University Extension and Outreach Online Store (ISU Extension publication **IPM 52**).

Results show double benefits of growing SCN-resistant soybeans

Although the performance of individual soybean varieties varied, the SCN-resistant soybean varieties had greater yields (figure 3) and consistently lower SCN soil egg population densities (figure 4) than the susceptible varieties in the experiments in 2015. Many individual SCN-resistant soybean varieties had greater yields than the mean yields shown in figure 3.




Figure 3. Mean yields of all SCN-resistant soybean varieties compared to susceptible soybean varieties at the nine field experiments conducted in 2015.




Figure 4. Mean end-of-season or final SCN egg population densities for all SCN-resistant soybean varieties compared to susceptible soybean varieties at the nine field experiments conducted in 2015.

The weather during the 2015 growing season was not such that typically results in a large amount of damage from SCN, relative to the damage it causes in hot, dry years. But nonetheless, yields and SCN numbers showed the benefits of resistant varieties in helping farmers produce higher yields while keeping the SCN numbers in check.

Use additional tactics for integrated management of SCN

Although the results from the experiments we conducted in 2015 illustrate that resistant varieties yield more and allow less SCN reproduction than susceptible varieties, the effectiveness of SCN-resistant varieties is eroding. The reason for the loss of effectiveness is that almost all SCN-resistant soybean varieties in the Midwest possess resistance genes from a breeding line or source of resistance called PI 88788. And continued use of the same set of resistance genes has selected for SCN populations with the ability to reproduce on plants with those resistance genes. In other words, exposing natural populations of SCN to the same set of resistance genes from PI 88788 over time has allowed the very few nematodes in the field that originally could overcome PI 88788 resistance genes to build up in numbers. It's the same force of nature at play that resulted in the buildup of glyphosate-resistant weeds. An October 2015 ICM News [article](#) describes the current situation with SCN-resistant soybean varieties in greater detail.

And as with build up of weed resistance to herbicides, using multiple management strategies is the best approach to managing the problem of the build up of SCN on PI

88788 SCN resistance. Specifically, farmers should use all available SCN management tools to slow the build up of SCN on soybean varieties with PI 88788 resistance. Currently, other SCN management tactics include growing nonhost crops, such as corn, and using nematode-protectant seed treatments on seeds of resistant soybean varieties.

More information about SCN

More information about the biology and management of SCN is available at www.soybeancyst.info and soybeanresearchinfo.com. Iowa State University's management recommendations for SCN are available online in a downloadable format, [Soybean Cyst Nematode \(SCN\) Management Recommendations](#), IPM 63.

Summary

- Resistant soybean varieties are a key SCN management tool, but individual varieties vary in agronomic performance and SCN control.
- Visit www.isuscntrials.info or see ISU Extension publication IPM 52 for results of ISU evaluations of SCN-resistant varieties in 2015.
- SCN populations are building up (developing increased reproduction) on the common PI 88788 source of SCN resistance.
- SCN-resistant varieties produced higher yields and resulted in lower SCN numbers than susceptible varieties in field experiments in 2015.
- Long-term management of SCN will require using nonhost crops and nematode-protectant seed treatments to help bolster the effectiveness of SCN-resistant varieties.

~~~~~

### Usage and limitations:

*Links to this article are strongly encouraged, and this article may be republished without further permission if published as written and if credit is given to the author, Integrated Crop Management News, and Iowa State University. If this article is to be used in any other manner, permission from the author is required.*

*This article was originally published on January 16, 2016. The information contained within may not be the most current and accurate depending on when it is accessed.*

**Category:** Plant Diseases

**Crop:****Soybean****Tags:**

SCN

SCN resistance

SCN management

**Author:****Greg Tylka** *Professor*

Dr. Greg Tylka is a professor in the Department of Plant Pathology and Microbiology at Iowa State University with extension and research responsibilities for management of plant-parasitic nematodes. The focus of Dr. Tylka's research program at Iowa State University is primarily the soybean cyst n...

---

[Site Index](#) [Site map](#) [Policies](#)